

PECAN NUTS: Information on products registered under Act 36 of 1947 for specific use in South Africa



The information listed below is provided as a guideline to growers.

The provider cannot accept any liability for its accuracy and users who rely on this information do so at their own risk.

NEMATOCIDE

Target	Active ingredient	Trade name	Reg. no.	Company	Active quantity
Plant parasitic nematodes	furfural	Crop Guard	L6864	Illovo Sugar Limited	900 g/L

Application rates: Initial application: 1 application of 5.0ml/m² followed by 2 applications of 2.5ml/m² at 2 to 3 week intervals. Drip/Micro-jet application: Make up a 10% solution with water, prior to application of product. Apply to a field irrigated to at least 70% field capacity. Follow the applications immediately with a minimum of 10 mm of water. Area to be treated (number of m²). Treat the tree basin, under the tree canopy for an overall surface spray or the area covered by the drip or micro-irrigation system if applied through these. Notes: Applications must be undertaken during the spring and autumn root flush periods. In instances of high nematode infestation, a further application can be made during the 3rd root flush period. When applied through the irrigation system, it will only control nematodes present in the area reached by the dripper or micro-jet.

Target	Active ingredient	Trade name	Reg. no.	Company	Active quantity
Plant parasitic nematodes	fenamiphos	Nemacur 100 GR	L2056	Villa Crop Protection (Pty) Ltd	100 g/kg

Application rates: 10 g/m² basin area. Apply the granules evenly, in Spring, to a trash free soil surface in the tree basin area. Irrigate immediately after application with at least 25 mm water. 120 day withholding period.